


## IMAGES IN CARDIOLOGY

# Aortopulmonary Artery Dissection

Hideki Itoh, MD,\* Takashi Yamamoto, MD,\* Hiroyuki Sugihara, MD,† Takao Saotome, MD,‡  
Yutaka Eguchi, MD,‡ Tohru Asai, MD,§ Minoru Horie, MD\*

*Shiga, Japan*



From the Departments of  
\*Cardiovascular and  
Respiratory Medicine,  
†Pathology, ‡Critical and  
Intensive Care Medicine,  
and §Surgery, Shiga  
University of Medical  
Science, Shiga, Japan.  
Manuscript received  
February 13, 2009,  
accepted February 18, 2009.

**A** 71-year-old man was brought to the hospital by ambulance and admitted because of intermittent back pain. Physical examination revealed continuous heart murmur. Chest X-ray showed cardiomegaly and pulmonary congestion. The patient received continuous hydration for renal dysfunction and severe metabolic acidosis (base excess of  $-16.5$  mmol/l and pH of 7.24) and underwent enhanced chest computed tomography. The chest computed tomography showed not only ascending aortic dissection of Stanford type A but also pulmonary artery dissection with an aortopulmonary window (**red arrow** = aortopulmonary window, T = true lumen, F = false lumen). An aortopulmonary shunt could not only increase pulmonary circulation and cause untreatable congestive pulmonary edema but also cause severe metabolic acidosis. Aortopulmonary artery dissection is a very rare disease but is fatal, requiring the surgical repair as rapidly as possible.